

A successful surgery requires a partnership between you and Dr. Butterfield

The following instructions are essential to a safe experience and good outcome. Use this as a checklist as you approach your surgery date. If you are unable to comply with these instructions, you must notify our office as soon as possible. As a result, your surgery may have to be postponed or delayed, at the judgment of Dr. Butterfield. This is essential to your health and safety.

### THREE WEEKS OR MORE BEFORE SURGERY

There may be several weeks between your decision to have surgery and your actual surgical date. During this time there are several important considerations:

**Practice proper fitness:** You need not engage in an aggressive or new fitness routine; however practicing good fitness habits is an important factor in your overall health and well-being. Stretching exercises and low-weight strength training now, can help to enhance your posture and your strength in the weeks following surgery.

**Good nutrition.** Eat well during the weeks prior to surgery. Crash dieting, over-eating or high alcohol intake can greatly affect your overall health and well-being. A healthy, balanced diet is essential. Also, begin taking a multi-vitamin daily.

**Stop smoking.** Smoking can greatly impair your ability to heal. You must be nicotine and smoke-free for at least 6 weeks prior to surgery. You must also be free of any nicotine patch or nicotine-based products for a minimum of 6 weeks prior to surgery.

**Lead a healthy lifestyle.** In the weeks prior to surgery maintain the best of health and hygiene. A lingering cold, virus or other illness can result in your surgery being rescheduled. Make certain to address any illness immediately, and advise our office of any serious illness or change in your health.

**Prepare and plan.** Schedule any time off of work, and any support you will need at home in the days following surgery, including housework, childcare, shopping and driving. Make certain a responsible adult is enlisted and confirmed to drive you to and from surgery, and that someone is confirmed available to stay with you around the clock for 24 hours, following surgery.

**Relax and enjoy life.** Stress and anxiety over life's daily events, and even your planned surgery can affect you. While some anxiety is common, any serious stress, or distress over the thought of surgery is something you must discuss with our office. We are here to support you and answer all of your questions. We want your decision to be one made with confidence.

**TWO to THREE WEEKS BEFORE SURGERY**

This is an important planning and preparation time. Follow all of the skincare and health habits you have begun in addition to the following:

\_\_\_\_\_ **Prepare and plan:** Put your schedule together for the day before, day of and first few days following the procedure. Share this with all of your key support people.

\_\_\_\_\_ **STOP taking the following for the duration before your surgery. Taking any of the following can increase your risk of bleeding and other complications:**

- |  |  |
|--|--|
| <input type="checkbox"/> Aspirin and medications containing aspirin          | <input type="checkbox"/> Garlic Supplements              |
| <input type="checkbox"/> Ibuprofen and anti-inflammatory agents (all NSAIDs) | <input type="checkbox"/> Green Tea or green tea extracts |
| <input type="checkbox"/> Vitamin E   | <input type="checkbox"/> St. John's Wort                 |
| <input type="checkbox"/> Coumadin/Warfarin (discuss with office)             | <input type="checkbox"/> Estrogen supplements            |
|  | <input type="checkbox"/> All other medications indicated |

\_\_\_\_\_ **Pre-operative clearance and information:** The pre-admission testing office of the hospital will contact you between 2 and 10 days prior to your operation. They will offer you the choice of having a history and physical performed by the hospital staff or by your primary care physician. This choice is entirely up to you. If your PCP completes the history and physical, the paperwork should be faxed to the hospital, as well as any laboratory tests. The history and physical form is included in the pre-operative packet. Simply give this paperwork to your PCP to complete.

\_\_\_\_\_ **Fitness:** Don't over-do it. Avoid anything strenuous or that could potentially cause injury.

\_\_\_\_\_ **Good nutrition:** Continue taking your supplements as directed.

\_\_\_\_\_ **NO SMOKING:** Stay away from second-hand smoke, too. Your healing and health depend heavily on this.

\_\_\_\_\_ **Avoid sun exposure:** Sun damaged skin can more readily produce irregular scars.

**ONE WEEK BEFORE SURGERY**

\_\_\_\_\_ **Confirm your day of surgery plans.** This includes your transportation and after-care (a responsible adult for the first 24 hours, around the clock).

**Pre-operative shopping list**

\_\_\_\_\_ The following is a list of items that should be purchased prior to surgery in order to prepare for and more easily recover from surgery.

- |  |  |
|--|--|
| <input type="checkbox"/> <b>Prescriptions</b> (have your prescriptions filled prior to surgery to save time on the way home when you won't be feeling up to stopping).                       | <input type="checkbox"/> <b>Germ-inhibiting soap</b> , such as Dial, Safeguard, or Lever 2000 (to bath with prior to surgery in order to minimize germs).  |
| <input type="checkbox"/> <b>Tylenol</b> (or a generic form of this drug) This will be the drug of choice once you do not need the prescription strength pain medications.                    | <input type="checkbox"/> <b>Frozen peas</b> (only for those patient undergoing facial procedures). These are great as "ice packs" for facial areas. Get 2-4 packages so that you can use 1 or 2 and have the others freezing |
| <input type="checkbox"/> <b>Multivitamin</b> (to take prior to surgery and during your recovery for maximum health).   | <input type="checkbox"/> <b>Bacitracin ointment</b> and <b>4x4 gauze</b> (to cover incisions with for the first week or so).   |
| <input type="checkbox"/> <b>Hydrogen peroxide</b> and <b>Q-tips</b> (to clean around drains).  | <input type="checkbox"/> <b>Stool softener</b> (e.g. Colace) and <b>laxative</b> (e.g. Dulcolax).  |
| <input type="checkbox"/> <b>Consider renting an electric lift chair</b> , if you are having a tummy tuck, buttock lift, or lower body lift. The office will be happy to assist you with this |  |

\_\_\_\_\_ **Continue to practice healthy habits**, nutrition and fitness. No strenuous exercise. No saunas, hot tubs, steam baths or mud wraps. **No smoking.**

\_\_\_\_\_ **Find your comfort zone.** Locate the most comfortable place where you can gently recline and recover. You don't want to be testing locations or pillows the day of surgery. Shop for magazines, books and other things to keep you busy and entertained in the day or two following surgery.

\_\_\_\_\_ **Relax.** Call our office with any unusual anxiety or concerns. Get plenty of rest. If you have trouble sleeping, call our office.

**ONE DAY BEFORE SURGERY**

\_\_\_\_\_ **Pack your bag for the day of surgery.** This should include:

- |   |  |
|---|--|
| <input type="checkbox"/> All paperwork                          | <input type="checkbox"/> Reading Glasses                   |
| <input type="checkbox"/> Your identification                    | <input type="checkbox"/> Chapstick                         |
| <input type="checkbox"/> All prescription medications           | <input type="checkbox"/> Saltines in case of nausea during |
| <input type="checkbox"/> Your post-surgical compression garment | your ride home   |

\_\_\_\_\_ **Expect a pre-anesthesia call to review your state of health for surgery**

\_\_\_\_\_ **Confirm your route to and from surgery or the recovery center, with the responsible adult who will drive you.** Also confirm plans with your 24-hour support person and make certain he or she has all of your post-operative instructions.

\_\_\_\_\_ **Shower as directed.** Use an anti-bacterial, fragrance-free soap, such as Lever 2000 or dial. Shampoo your hair. Do not use any hair gel or other styling products, scented skin creams or moisturizers. Do not use any deodorant, hair spray, perfume or cosmetics. Remove all finger nail and toe nail polish.

\_\_\_\_\_ **Wax or shave.** It may be uncomfortable to do so in the days immediately after surgery.

\_\_\_\_\_ **Do not eat or drink anything after 12 pm.** No candy, gum or mints. Anything more than a small amount of water as needed for brushing teeth or swallowing medication may result in the need to cancel surgery.

\_\_\_\_\_ **RELAX!** Get plenty of rest and avoid unnecessary stress.

**THE DAY OF SURGERY**

\_\_\_\_\_ **NOTHING by mouth.** Anything more than a small amount of water as needed for brushing teeth or swallowing medication may result in the need to cancel surgery. This includes candy, gum, mints.

\_\_\_\_\_ **Dress appropriately.**

- **Do not wear cosmetics, jewelry of any kind, contact lenses, hair clips, body piercing:** (If there is something you cannot remove, let the admitting nurse know right away.)
- **Wear comfortable, clean, loose-fitting clothing:** Do not wear jeans or any tight-fitting bottom; rather have a pair of loose, drawstring sweatpants to wear home. Wear slip on, flat shoes with a slip proof sole; no heels. Wear clean cotton socks, as the operating room can feel cool. For your comfort, wear a zip or button front top. No turtlenecks.

I look forward to seeing you prior to surgery in the same day surgery area of the hospital. If you have any last minute questions, we will have time to discuss them. I will also be marking you for surgery at that time.

Date of Birth: \_\_\_\_\_

Date to be admitted: \_\_\_\_\_

Reason for procedure: \_\_\_\_\_

**History of Present Illness:**

**Past History (Psycho/Social History):**

**Drug or Other Significant Allergies:**

**CURRENT MEDICATIONS (including ASA):**

All "Yes" answers require amplification or comment

	YES	NO	Comment
Diabetes			
History of Steroid Treatment			
Bleeding Tendency			
Weight Loss			
Smoking:			
Currently smoking			
History of smoking			
Review of Systems:			
Pain or discomfort			
Cardiovascular:			
Chest Pain			
History of MI			
Syncope			
Other Pertinent symptoms			
Respiratory:			
Effort Intolerance			
History of Asthma			
Cough			
Other Pertinent symptoms			
Cardiovascular:			
Chest Pain			
History of MI			
Syncope			
Other Pertinent symptoms			
Neurological:			
History of Transient neurological symptoms			
Other Pertinent symptoms			
Gastrointestinal:			
Abdominal Pain			
Nausea			
History of Hepatitis			
Other Pertinent symptoms			
Reproductive:			
Other Pertinent symptoms			Last Menstrual Period Date:
Other: (musculoskeletal, endocrine, GU, etc.)			

**Physical Examination**

--

<b>GENERAL APPEARANCE:</b>	<b>BP:</b>		
	<b>PULSE:</b>		
	<b>RESP:</b>		
	<b>TEMP:</b>		
<b>HEENT:</b>			
<b>NECK:</b>		<b>YES</b>	<b>NO</b>
	<b>Bruits</b>		
<b>CHEST &amp; LUNGS</b>			
<b>BREASTS:</b>			
<b>HEART:</b>		<b>YES</b>	<b>NO</b>
	<b>Murmur</b>		
<b>ABDOMEN:</b>			
<b>PELVIC/RECTAL INGUINO-GENITAL:</b>			
<b>EXTREMITIES:</b>		<b>YES</b>	<b>NO</b>
	<b>Venous Stasis</b>		
<b>NEUROLOGICAL:</b>			
<b>ASSESSMENT/CONCLUSION:</b>			

**RISK ASSESSMENT: ASA CRITERIA (Circle one)**

1. Normal healthy patient      2. Mild disease, no functional impairment      3. Severe systemic disease  
4. Disease is constant threat to life      5. Moribund, not expected to recover

**Signature:** \_\_\_\_\_ **Name Printed:** \_\_\_\_\_  
**Date of Examination:** \_\_\_\_\_ **Phone Number:** \_\_\_\_\_

**Re-assessment: Patient identified, chart reviewed, condition unchanged. Appropriate to proceed with planned sedation.**  
Physician initials \_\_\_\_\_ Date: \_\_\_\_\_

**THE JEWISH HOSPITAL PREADMISSION TESTING PROTOCOLS**

TEST	CRITERIA
<b>History and Physical</b>	<b>All patients need a current health screening updated within 30 days</b>
<b>Hemoglobin</b>	<b>History of Anemia History of bleeding, i.e.: Hematuria, vaginal bleeding All patients who receive type and screens</b>
<b>Basic Metabolic Panel</b>	<b>Patients with renal disease. (BMP as close to day of surgery as possible) Patients taking steroids. (BMP within 30 days)</b>
<b>Blood Sugar</b>	<b>Patients who are diabetic</b>
<b>Potassium Level</b>	<b>Patients taking potassium depleting diuretics *see listing below Patients taking digitalis</b>
<b>Coag profile</b>	<b>Patients recently undergoing chemotherapy Hepatic Disease Bleeding Disorder Anticoagulant therapy taken within the last three days.</b>
<b>Blood levels of therapeutic drugs</b>	<b>Digitalis Lithium Seizure medications, ie: tegretol, dilantin, Phenobarbital Theophylline, depakene (if using for a seizure disorder)</b>
<b>EKG – if EKG available and patient’s health status has not changed since reading, do not repeat</b>	<b>Patients with cardiac and peripheral vascular disease, history of dysrhythmias, hypertension, MVP, Graves Disease, Diabetes Morbid obesity (BMI &gt; 40) Thoracotomy surgery. Surgery planned for greater than 3 hours.</b>
<b>CXR</b>	<b>Patients with acute pulmonary symptoms.</b>
<b>Urine Pregnancy test</b>	<b>Performed on day of surgery for females with onset of menses up to one year post menses.</b>
<b>Type and Screen</b>	<b>According to blood bank protocol.</b>

**DIURECTICS THAT REQUIRE SERUM POTASSIUM LEVELS**

**Thiazide and Thiazide-like Diurectics**

**DIURIL – chlorothiazide  
HYDRODIURIL – hydrochlorothiazide  
NATURETIN – bendroflumethiazide  
ENDURON – methyclothiazide  
EXNA – benzthiazide**

**METAHYDRIN - trichlormethiazide  
RENESE - polythiazide  
HYDROMOX - quinethazone  
ZAROXOLYN - metolazone  
HYGROTON - chlorthalidone  
DIUCARDIN - hydroflumethiazide**

**Loop diurectics**

**LASIX – furosemide  
BUMEX – bumetanide  
EDECIN – ethacrynic acid  
DEMEDEX - torsemide**

THE CHRIST HOSPITAL  
CINCINNATI, OHIO 45219  
HISTORY AND PHYSICAL EXAMINATION  
R-54A REV. 7/99  
PAGE 1 OF 2

ADMITTING/TESTING FAX # 585-1273

DATE OF EXAM \_\_\_\_\_

REASON FOR ADMISSION/INDICATION FOR PROCEDURE: \_\_\_\_\_

HISTORY OF PRESENT ILLNESS: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

DRUG OR OTHER SIGNIFICANT ALLERGIES: \_\_\_\_\_

FAMILY AND SOCIAL HISTORY: \_\_\_\_\_

PAST HISTORY: \_\_\_\_\_

DIABETES: no \_\_\_ yes \_\_\_; Hx. Steroid Rx: no \_\_\_ yes \_\_\_; Hx. of Diuretic Rx: no \_\_\_ yes \_\_\_

Bleeding Tendency: no \_\_\_ yes \_\_\_

CURRENT MEDICATIONS: \_\_\_\_\_

\_\_\_\_\_

ALL YES ANSWERS REQUIRE COMMENT:

R.O.S: Pain or Discomfort no \_\_\_ yes \_\_\_ specify

Weight loss: no \_\_\_ yes \_\_\_

CARDIOVASCULAR:

Chest pain no \_\_\_ yes \_\_\_

Hx of MI no \_\_\_ yes \_\_\_

Syncope no \_\_\_ yes \_\_\_

Hx. of Deep Vein Thrombosis no \_\_\_ yes \_\_\_

Other pertinent sx. no \_\_\_ yes \_\_\_

RESPIRATORY:

Hx of Asthma no \_\_\_ yes \_\_\_

Cough no \_\_\_ yes \_\_\_

Smoke no \_\_\_ yes \_\_\_

Other pertinent sx. no \_\_\_ yes \_\_\_

NEUROLOGICAL:

Hx. of transient neurological sx. no \_\_\_ yes \_\_\_

Other pertinent symptoms no \_\_\_ yes \_\_\_

RENAL:

HX: Kidney or bladder disease no \_\_\_ yes \_\_\_

Other pertinent symptoms no \_\_\_ yes \_\_\_

GASTROINTESTINAL:

Abdominal pain no \_\_\_ yes \_\_\_

Nausea no \_\_\_ yes \_\_\_

Hx of Hepatitis no \_\_\_ yes \_\_\_

Alcohol use no \_\_\_ yes \_\_\_

Other pertinent sx: no \_\_\_ yes \_\_\_

REPRODUCTIVE:

Last Menstrual Period Date:

OTHER: (Musculoskeletal, endocrine, GU etc) \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

PHYSICAL EXAMINATION:

TEMP ____	PR ____	RESP ____	BP ____
-----------	---------	-----------	---------

GENERAL APPEARANCE:

MENTAL STATUS:

HEENT:

NECK:

Bruits: no \_\_\_\_ yes \_\_\_\_

CHEST AND LUNGS:

Breasts

HEART:

Murmur: no \_\_\_\_ yes \_\_\_\_

ABDOMEN:

PELVIC/RECTAL/INGUINO GENITAL:

EXTREMITIES:

Venous Stasis no \_\_\_\_ yes \_\_\_\_

NEUROLOGICAL:

DIAGNOSIS \_\_\_\_\_

ASSESSMENT/PLAN:

SIGNATURE: \_\_\_\_\_ NAME PRINTED: \_\_\_\_\_

# **INFORMED CONSENT – BREAST RECONSTRUCTION WITH LATISSIMUS MUSCLE FLAP**

## **INSTRUCTIONS**

This informed-consent document has been prepared to help inform you of breast reconstruction with latissimus muscle flap surgery, its risks, and alternative treatments.

It is important that you read this information carefully and completely. Please initial each page, indicating that you have read the page and sign the consent for surgery as proposed by your plastic surgeon and agreed upon by you.

## **GENERAL INFORMATION**

There is a variety of surgical techniques for breast reconstruction. Most mastectomy patients are medically appropriate for breast reconstruction, either immediately following breast removal or at a later time. The best candidates, however, are women whose cancer, as far as can be determined, seems to be eliminated by mastectomy. There are legitimate reasons to delay breast reconstruction. Some women may be advised by their surgeon or oncologist to wait until other forms of necessary cancer treatment are completed. Other patients may require more complex breast reconstruction procedures. Women who smoke, or who have other health conditions, such as obesity or high blood pressure, may be advised to postpone surgery. In any case, being informed of your options concerning breast reconstruction can help you prepare for a mastectomy with a more positive outlook on the future.

Breast reconstruction has no known effect on altering the natural history of breast cancer or interfering with other forms of breast cancer treatment such as chemotherapy or radiation.

In breast reconstruction with the latissimus dorsi, a muscle located on the back along with its attached skin is transferred to the chest region for the breast reconstruction procedure. The muscle flap maintains its own blood supply, and helps nourish the tissue that is transferred to the chest wall region. There are several variations on the surgical technique of latissimus muscle flap breast reconstruction, including microvascular surgery to attach the flap to the chest region. In some cases, your plastic surgeon may recommend that a breast implant be inserted underneath the muscle flap to give the breast mound additional projection.

Muscle flap techniques of breast reconstruction are useful in the following situations:

- Inadequate chest wall tissue for breast reconstruction with implants or expanders
- Past history of radiation to chest wall after mastectomy
- Patient with concerns about silicone breast implant/expander
- Failure of earlier breast reconstruction

**Contraindications to latissimus muscle flap breast reconstruction procedure exist:**

- A patient who is medically or psychologically unsuitable for breast reconstruction
- Previous injury to the latissimus dorsi muscle from surgery or other treatments

**A separate consent form for the use of breast implants in conjunction with breast reconstruction with latissimus muscle flap is necessary.**

## **ALTERNATIVE TREATMENTS**

Latissimus muscle flap breast reconstruction is an elective surgical operation. Alternative treatment would consist of the use of external breast prostheses or padding, tissue expansion technique of breast reconstruction, breast implants, or the transfer of other body tissues for breast reconstruction. Risks and potential complications are associated with alternative surgical forms of treatment.

## **INFORMED CONSENT – BREAST RECONSTRUCTION WITH LATISSIMUS MUSCLE FLAP**

### **RISKS OF BREAST RECONSTRUCTION WITH LATISSIMUS MUSCLE FLAP**

Every surgical procedure involves a certain amount of risk, and it is important that you understand the risks involved with breast reconstruction with latissimus muscle flap and the possible use of a breast implant in addition to the muscle flap. In the event that a latissimus muscle flap is used without a breast implant, risks associated with breast implants would not be applicable. An individual's choice to undergo a surgical procedure is based on the comparison of the risk to potential benefit. Although the majority of women do not experience the following complications, you should discuss each of them with your plastic surgeon to make sure you understand the risks, potential complications, and consequences of breast reconstruction with latissimus muscle flap.

**Bleeding-** It is possible, though unusual, to experience a bleeding episode during or after surgery. Should post-operative bleeding occur, it may require emergency treatment to drain accumulated blood (hematoma) or blood transfusion. Intra-operative blood transfusion may also be required. Do not take any aspirin or anti-inflammatory medications for ten days before or after surgery, as this may increase the risk of bleeding. Non-prescription "herbs" and dietary supplements can increase the risk of surgical bleeding. Hematoma can occur at any time following injury to the breast. If blood transfusions are necessary to treat blood loss, there is the risk of blood-related infections such as hepatitis and HIV (AIDS). Heparin medications that are used to prevent blood clots in veins can produce bleeding and decreased blood platelets.

**Infection-** An infection is unusual after this type of surgery. Should an infection occur, treatment including antibiotics, hospitalization, or additional surgery may be necessary. Infections with the presence of a breast implant (if used) are harder to treat than infections in normal body tissues. If an infection does not respond to antibiotics, the breast implant may have to be removed. After the infection is treated, a new breast implant can usually be reinserted.

**Change in Nipple and Skin Sensation-** Breast reconstruction cannot restore normal sensation to your breast or nipple. Skin that is transferred as part of the muscle flap will lack sensation. Numbness may occur in the skin on the back where the latissimus muscle was located. Changes in sensation may affect sexual response or the ability to breast-feed a baby.

**Skin Contour Irregularities-** Contour and shape irregularities may occur. One breast may be smaller than the other. Nipple position and shape will not be identical one side to the next. Residual skin irregularities at the ends of the incisions or "dog ears" are always a possibility when there is excessive redundant skin. This may improve with time, or it can be surgically corrected.

**Sutures-** Most surgical techniques use deep sutures. You may notice these sutures after your surgery. Sutures may spontaneously poke through the skin, become visible or produce irritation that requires suture removal.

**Skin Discoloration / Swelling-** Some bruising and swelling normally occurs following a breast reconstruction. The skin in or near the surgical site can appear either lighter or darker than surrounding skin. Although uncommon, swelling and skin discoloration may persist for long periods of time and, in rare situations, may be permanent. Additionally, these areas may have exaggerated responses to hot or cold temperatures.

**Scarring-** All surgery leaves scars, some more visible than others. Although good wound healing after a surgical procedure is expected, abnormal scars may occur within the skin and deeper tissues. Scars may be unattractive and of different color than the surrounding skin tone. Scar appearance may also vary within the same scar. Scars may be asymmetrical (appear different on the right and left side of the body). There is the possibility of visible marks in the skin from sutures. In some cases, scars may require surgical revision or treatment.

## **INFORMED CONSENT – BREAST RECONSTRUCTION WITH LATISSIMUS MUSCLE FLAP**

**Delayed Healing and Loss of Flap-** Wound disruption or delayed wound healing is possible. It is possible to have areas of the chest wall or latissimus dorsi muscle flap die. This may require frequent dressing changes or further surgery to remove the non-living tissue. Individuals who have decreased blood supply to breast tissue from past surgery or radiation therapy may be at increased risk for delayed wound healing and poor surgical outcome. **Smokers have a greater risk of skin loss and wound healing complications.**

**Damage to Deeper Structures-** There is the potential for injury to deeper structures including nerves, blood vessels, muscles, and lungs (pneumothorax) during any surgical procedure. The potential for this to occur varies according to the type of procedure being performed. Injury to deeper structures may be temporary or permanent.

**Fat Necrosis-** Fatty tissue found in the flap or skin may die. This may produce areas of firmness within the skin. Additional surgery to remove areas of fat necrosis may be necessary (breast biopsy). There is the possibility of contour irregularities in the skin that may result from fat necrosis.

**Seroma-** Pockets of tissue fluid sometimes develop either in the back or in the chest wall after a latissimus muscle flap breast reconstruction. Additional procedures to drain this fluid accumulation may be necessary.

**Breast Implants-** Risks associated with the potential use of breast implants are covered in a separate informed-consent form.

**Implant Extrusion-** Lack of adequate tissue coverage may result in exposure and extrusion of a breast implant, if used, in addition to the latissimus muscle flap. If tissue breakdown occurs and the breast implant becomes exposed, removal is necessary.

**Firmness-** Excessive firmness of the breast can occur after surgery due to internal scarring or scarring around a breast implant if one is used. The occurrence of this is not predictable and additional treatment or surgery may be necessary. Radiation therapy to the chest region after breast reconstruction with a latissimus muscle flap may produce unacceptable firmness or other long-term complications.

**Asymmetry-** Some breast asymmetry naturally occurs in most women. Differences in breast and nipple shape, size, or symmetry may also occur after surgery. Additional surgery may be necessary to correct asymmetry after a breast reconstruction with latissimus muscle flap.

**Allergic Reactions-** In rare cases, local allergies to tape, suture material and glues, blood products, topical preparations or injected agents have been reported. Serious systemic reactions including shock (anaphylaxis) may occur in response to drugs used during surgery and prescription medicines. Allergic reactions may require additional treatment.

**Surgical Anesthesia-** Both local and general anesthesia involve risk. There is the possibility of complications, injury, and even death from all forms of surgical anesthesia or sedation.

**Unsatisfactory Result-** Although good results are expected, there is no guarantee or warranty expressed or implied, on the results that may be obtained. You may be disappointed with the results breast reconstruction surgery. Asymmetry in muscle flap placement, unanticipated breast shape and size, loss of function, wound disruption, poor healing, and loss of sensation may occur after surgery. Breast size may be incorrect. Unsatisfactory surgical scar location or visible deformities at the ends of the incisions (dog ears) may occur. Breast reconstruction by any technique may fail due to complications attributable to the mastectomy surgery or from chemotherapy/radiation therapy treatments that are independent of the latissimus muscle flap procedure. It may be necessary to perform additional surgery to improve your results.

**Loss of Latissimus Muscle Function-** There is anticipated loss of normal function in the latissimus dorsi muscle after it is transferred to the chest wall. Weakness in movements of the shoulder and upper arm can occur.

## **INFORMED CONSENT – BREAST RECONSTRUCTION WITH LATISSIMUS MUSCLE FLAP**

**Shock-** In rare circumstances, your surgical procedure can cause severe trauma, particularly when multiple or extensive procedures are performed. Although serious complications are infrequent, infections or excessive fluid loss can lead to severe illness and even death. If surgical shock occurs, hospitalization and additional treatment would be necessary.

**Pain-** You will experience pain after your surgery. Pain of varying intensity and duration may occur and persist after surgery. Chronic pain may occur very infrequently from nerves becoming trapped in scar tissue after a breast reconstruction.

**Thrombosed Veins-** Thrombosed veins, which resemble cords, occasionally develop in the area of the breast and resolve without medical or surgical treatment.

**Cardiac and Pulmonary Complications-** Surgery, especially longer procedures, may be associated with the formation of, or increase in, blood clots in the venous system. Pulmonary complications may occur secondarily to both blood clots (pulmonary emboli), fat deposits (fat emboli) or partial collapse of the lungs after general anesthesia. Pulmonary emboli can be life threatening or fatal in some circumstances. Inactivity and other conditions may increase the incidence of blood clots traveling to the lungs causing a major blood clot that may result in death. It is important to discuss with your physician any past history of blood clots or swollen legs that may contribute to this condition. Cardiac complications are a risk with any surgery and anesthesia, even in patients without symptoms. If you experience shortness of breath, chest pains, or unusual heart beats, seek medical attention immediately. Should any of these complications occur, you may require hospitalization and additional treatment.

### **ADDITIONAL ADVISORIES**

**Breast Disease-** Current medical information does not demonstrate an increased risk of breast disease, breast cancer, or recurrence of breast cancer in women who have reconstructive breast surgery. Breast disease and breast cancer can occur independently of breast surgery. Individuals with a personal history or family history of breast cancer may be at a higher risk of developing breast cancer than a woman with no family history of this disease. It is recommended that all women perform regular self-examination of their breasts, have mammograms according to American Cancer Society guidelines, and seek professional care should a breast lump be detected.

**Interference with Sentinel Lymph Node Mapping Procedures-** Breast surgery procedures that involve cutting through breast tissue, similar to a breast biopsy, can potentially interfere with diagnostic procedures to determine lymph node drainage of breast tissue to stage breast cancer.

**Breast and Nipple Piercing Procedures-** Individuals who currently wear body-piercing jewelry in the breast region are advised that a breast infection could develop from this activity.

**Pregnancy and Breast Feeding-** There is no evidence that muscle flap surgery has any effect on fertility or pregnancy. If a woman has undergone a mastectomy, it is unlikely that she would be able to breast feed a baby on the affected side.

**Long-Term Results-** Subsequent alterations in breast shape may occur as the result of aging, sun exposure, weight loss, weight gain, pregnancy, menopause, or other circumstances not related to your surgery. Breast sagginess may normally occur.

**Female Patient Information-** It is important to inform your plastic surgeon if you use birth control pills, estrogen replacement, or if you suspect you may be pregnant. Many medications including antibiotics may neutralize the preventive effect of birth control pills, allowing for conception and pregnancy.

**Intimate Relations After Surgery-** Surgery involves coagulating of blood vessels and increased activity of any kind may open these vessels leading to a bleed, or hematoma. Activity that increases your pulse or heart rate may cause additional bruising, swelling, and the need for return to surgery and control of bleeding. It is wise to refrain from sexual activity until your physician states it is safe.

## **INFORMED CONSENT – BREAST RECONSTRUCTION WITH LATISSIMUS MUSCLE FLAP**

### **Smoking, Second-Hand Smoke Exposure, Nicotine Products (Patch, Gum, Nasal Spray)-**

Patients who are currently smoking, use tobacco products, or nicotine products (patch, gum, or nasal spray) are at a greater risk for significant surgical complications of skin dying, delayed healing, and additional scarring. Individuals exposed to second-hand smoke are also at potential risk for similar complications attributable to nicotine exposure. Additionally, smoking may have a significant negative effect on anesthesia and recovery from anesthesia, with coughing and possibly increased bleeding. Individuals who are not exposed to tobacco smoke or nicotine-containing products have a significantly lower risk of this type of complication. Please indicate your current status regarding these items below:

\_\_\_\_\_ I am a non-smoker and do not use nicotine products. I understand the risk of second-hand smoke exposure causing surgical complications.

\_\_\_\_\_ I am a smoker or use tobacco / nicotine products. I understand the risk of surgical complications due to smoking or use of nicotine products.

It is important to refrain from smoking at least 6 weeks before surgery and until your physician states it is safe to return, if desired.

**Medications-** There are potential adverse reactions that occur as the result of taking over-the-counter, herbal, and/or prescription medications. Be sure to check with your physician about any drug interactions that may exist with medications which you are already taking. If you have an adverse reaction, stop the drugs immediately and call your plastic surgeon for further instructions. If the reaction is severe, go immediately to the nearest emergency room. When taking the prescribed pain medications after surgery, realize that they can affect your thought process and coordination. Do not drive, do not operate complex equipment, do not make any important decisions and do not drink any alcohol while taking these medications. Be sure to take your prescribed medication only as directed.

**Mental Health Disorders and Elective Surgery-** It is important that all patients seeking to undergo elective surgery have realistic expectations that focus on improvement rather than perfection. Complications or less than satisfactory results are sometimes unavoidable, may require additional surgery and often are stressful. Please openly discuss with your surgeon, prior to surgery, any history that you may have of significant emotional depression or mental health disorders. Although many individuals may benefit psychologically from the results of elective surgery, effects on mental health cannot be accurately predicted.

### **ADDITIONAL SURGERY NECESSARY (Re-Operations)**

Many variable conditions may influence the long-term result of breast reconstruction with latissimus muscle flap surgery. Secondary surgery may be necessary to perform additional tightening or repositioning of the breasts. Should complications occur, additional surgery or other treatments may be necessary. Even though risks and complications occur infrequently, the risks cited are the ones that are particularly associated with breast reconstruction with latissimus muscle flap surgery. Other complications and risks can occur but are even more uncommon. The practice of medicine and surgery is not an exact science. Although good results are expected, there is no guarantee or warranty expressed or implied on the results that may be obtained. In some situations, it may not be possible to achieve optimal results with a single surgical procedure.

## **INFORMED CONSENT – BREAST RECONSTRUCTION WITH LATISSIMUS MUSCLE FLAP**

### **PATIENT COMPLIANCE**

Follow all physician instructions carefully; this is essential for the success of your outcome. It is important that the surgical incisions are not subjected to excessive force, swelling, abrasion, or motion during the time of healing. Personal and vocational activity needs to be restricted. Protective dressings and drains should not be removed unless instructed by your plastic surgeon. Successful post-operative function depends on both surgery and subsequent care. Physical activity that increases your pulse or heart rate may cause bruising, swelling, fluid accumulation and the need for return to surgery. It is wise to refrain from intimate physical activities after surgery until your physician states it is safe. It is important that you participate in follow-up care, return for aftercare, and promote your recovery after surgery.

### **HEALTH INSURANCE**

Most insurance carriers consider breast reconstruction surgery a covered benefit. However, there may be additional requirements. Please review your health insurance subscriber-information pamphlet, call your insurance company, and discuss this further with your plastic surgeon. **Many insurance plans exclude coverage for secondary or revisionary surgery.**

### **FINANCIAL RESPONSIBILITIES**

The cost of surgery involves several charges for the services provided. The total includes fees charged by your surgeon, the cost of surgical supplies, anesthesia, laboratory tests, and possible outpatient hospital charges, depending on where the surgery is performed. Depending on whether the cost of surgery is covered by an insurance plan, you will be responsible for necessary co-payments, deductibles, and charges not covered. The fees charged for this procedure do not include any potential future costs for additional procedures that you elect to have or require in order to revise, optimize, or complete your outcome. Additional costs may occur should complications develop from the surgery. Secondary surgery or hospital day-surgery charges involved with revision surgery will also be your responsibility. **In signing the consent for this surgery/procedure, you acknowledge that you have been informed about its risk and consequences and accept responsibility for the clinical decisions that were made along with the financial costs of all future treatments.**

### **DISCLAIMER**

Informed-consent documents are used to communicate information about the proposed surgical treatment of a disease or condition along with disclosure of risks and alternative forms of treatment(s), including no surgery. The informed-consent process attempts to define principles of risk disclosure that should generally meet the needs of most patients in most circumstances.

However, informed-consent documents should not be considered all inclusive in defining other methods of care and risks encountered. Your plastic surgeon may provide you with additional or different information that is based on all the facts in your particular case and the current state of medical knowledge.

Informed-consent documents are not intended to define or serve as the standard of medical care. Standards of medical care are determined on the basis of all of the facts involved in an individual case and are subject to change as scientific knowledge and technology advance and as practice patterns evolve.

**It is important that you read the above information carefully and have all of your questions answered before signing the consent on the next page.**

**INFORMED CONSENT – BREAST RECONSTRUCTION  
WITH LATISSIMUS MUSCLE FLAP**

**CONSENT FOR SURGERY / PROCEDURE or TREATMENT**

1. I hereby authorize Dr. Jennifer Butterfield and such assistants as may be selected to perform the following procedure or treatment:

**BREAST RECONSTRUCTION WITH LATISSIMUS DORSI MUSCLE FLAP**

I have received the following information sheet:

**INFORMED CONSENT - BREAST RECONSTRUCTION WITH LATISSIMUS DORSI MUSCLE FLAP**

2. I recognize that during the course of the operation and medical treatment or anesthesia, unforeseen conditions may necessitate different procedures than those above. I therefore authorize the above physician and assistants or designees to perform such other procedures that are in the exercise of his or her professional judgment necessary and desirable. The authority granted under this paragraph shall include all conditions that require treatment and are not known to my physician at the time the procedure is begun.
3. I consent to the administration of such anesthetics considered necessary or advisable. I understand that all forms of anesthesia involve risk and the possibility of complications, injury, and sometimes death.
4. I acknowledge that no guarantee or representation has been given by anyone as to the results that may be obtained.
5. I consent to be photographed or televised before, during, and after the operation(s) or procedure(s) to be performed, including appropriate portions of my body, for medical, scientific or educational purposes, provided my identity is not revealed by the pictures.
6. For purposes of advancing medical education, I consent to the admittance of observers to the operating room.
7. I consent to the disposal of any tissue, medical devices or body parts which may be removed.
8. I consent to the utilization of blood products should they be deemed necessary by my surgeon and/or his/her appointees, and I am aware that there are potential significant risks to my health with their utilization.
9. I authorize the release of my Social Security number to appropriate agencies for legal reporting and medical-device registration, if applicable.
10. I understand that the surgeons' fees are separate from the anesthesia and hospital charges, and the fees are agreeable to me. If a secondary procedure is necessary, further expenditure will be required.
11. I realize that not having the operation is an option.
12. IT HAS BEEN EXPLAINED TO ME IN A WAY THAT I UNDERSTAND:
- a. THE ABOVE TREATMENT OR PROCEDURE TO BE UNDERTAKEN
  - b. THERE MAY BE ALTERNATIVE PROCEDURES OR METHODS OF TREATMENT
  - c. THERE ARE RISKS TO THE PROCEDURE OR TREATMENT PROPOSED

I CONSENT TO THE TREATMENT OR PROCEDURE AND THE ABOVE LISTED ITEMS (1-12). I AM SATISFIED WITH THE EXPLANATION.

\_\_\_\_\_  
Patient or Person Authorized to Sign for Patient

Date \_\_\_\_\_ Witness \_\_\_\_\_

## Post-Surgical Instructions: Breast Reconstruction with Latissimus Flap

Once your surgery is completed, you must follow all the instructions given to you in order to heal properly and have good outcomes.

The following instructions are your obligation. Use this as a checklist of progress as you heal. Included are normal post-surgical experiences and key health considerations that may be a cause of concern.

### TYPICAL POST-OPERATIVE SYMPTOMS

Typical symptoms of breast reconstruction and signs to watch for following breast reconstruction with a latissimus flap and breast implant include the following:

**Tightness in the chest or back region and stiffness; Tingling, burning or intermittent shooting pain:** These are normal experiences as the skin, muscles and tissue stretch to accommodate your implants, and as sensory nerves heal. Pain medication and muscle relaxants will help you cope with any discomfort. If you have drains, you may experience additional localized discomfort. **Consistent sharp pain should be reported to our office immediately.**

**Shiny skin or any itchy feeling:** Swelling can cause the breasts' skin to appear shiny. As the healing process advances, you may also find a mild to severe itchy feeling of the breasts. An antihistamine like Benadryl can help to alleviate severe, constant itchiness. **If the skin becomes red and hot to the touch, contact our office immediately.**

**Asymmetry, the breasts look different, or heal differently.** Breasts may look or feel quite different from one another in the days following surgery. This is normal; no two breasts are perfectly symmetrical in nature or following breast reconstruction surgery.

**A sloshing sound or sensation:** This is not the result of your implant filler, but rather of air that is trapped in the implant pocket and fluid that may naturally accumulate. This is perfectly normal and will resolve within 2-4 weeks.

### OUR OFFICE SHOULD BE CONTACTED IMMEDIATELY IF YOU EXPERIENCE ANY OF THE FOLLOWING:

- A high fever, (over 101.5°) severe nausea and vomiting, continued dizziness or incoherent behavior, such as hallucinations.
- Any pain that cannot be controlled by your pain medication.
- Bright red skin that is hot to the touch.
- Excessive bleeding or fluid seeping through the incisions.
- A severely misshapen breast or bruising that is localized to one breast or region of the chest.

To alleviate any discomfort, and to reduce swelling, you may apply cool, not cold compresses to the treated region: your breast or back donor site. Do not apply ice or anything frozen directly to the skin. Apply cool compresses, for no longer than 20-minute intervals.

Do not apply heat or cold.



## Post-Surgical Instructions: Breast Reconstruction with Latissimus Flap



### FOUR TO TEN DAYS FOLLOWING SURGERY

During this time you will progress with each day that passes. Ease into your daily activities. You will receive clearance to begin driving 1 to 2 weeks after your procedure (you must be off of narcotics), and we will discuss your return to work at your first post-operative visit.

- **Continue to cleanse wounds as directed; you may shower.** Take a warm, not hot shower. Do not take a bath. Limit your shower to 10 minutes. Apply bacitracin ointment to your incisions (breast and back) 2 x daily then apply a gauze pad, and use the post-operative bra (supplied by the hospital) to hold the dressings in place. Apply a fragrance free moisturizer to breast and surrounding skin, however not on your incisions.
- **Take antibiotic medications and supplements as directed.** Take pain medication and muscle relaxants only as needed.
- **Continue your breast massage. Wear your bra around the clock.**
- **Maintain daily walking.** Walking is essential every day to prevent the formation of blood clots. In addition, you may begin stretching and range of motion exercises.
- **Maintain a healthy diet. Do not smoke. Do not consume alcohol.**

### TWO to FOUR WEEKS FOLLOWING SURGERY

As you resume your normal daily activities, you must continue proper care and healing.

- **Continue your breast massage and wound care as directed.** You no longer need to use gauze on the incisions, as the drainage has stopped. Continue to apply bacitracin ointment to the incisions 2 x daily. You may start gentle scar massage, several times daily.
- **You may wish to switch from prescriptive pain medication to acetaminophen.**
- **Ease into light weight-bearing exercise.** No tennis, golf, softball or other sports with similar swinging motions. Avoid aerobic exercise that may cause a lot of bounce.
- **Do not smoke.** While incisions may have sealed, smoking deprives your body of necessary oxygen that can result in poorly healed, wide, raised scars.
- **Continue to wear a proper support bra.** The bra you first wore following surgery may feel somewhat loose. You may replace it, however **no under wires for 6 weeks**. You may sleep without a bra; however a camisole with built-in shelf support can be comfortable and provides added support as you continue to heal.
- **You may sleep flat.** However do not sleep on your stomach. If you are a side sleeper, a soft pillow under your mid-back and shoulders may offer more comfort and support than a single pillow under your head.
- **Practice good sun protection.** Do not expose your breasts or back to direct sunlight. If you are outdoors, apply at least an SPF 30 to the chest area at least 30 minutes prior to sun exposure. Your chest region, breast skin and donor site are highly susceptible to sunburn or the formation of irregular, darkened pigmentation.

## Post-Surgical Instructions: Breast Reconstruction with Latissimus Flap



### SIX WEEKS FOLLOWING SURGERY

Healing will progress and your breasts will settle into a more final shape and position.

- **You may ease into your regular fitness routine.** However realize that your upper body may require some time to return to prior strength.
- **Continue your breast massage;** this is essential for as long as you plan to have breast implants.
- **You may resume wearing under wires,** although these are not necessary.
- **Discomfort or tightness and tingling will resolve.** Any lingering nipple sensitivity or lack of sensation should begin to greatly improve.
- **No need to resume smoking.** You have now gone 10 weeks (4 weeks prior to surgery and 6 weeks following) without a cigarette. For your long-term health, there is no need to resume smoking.
- **We will discuss the next two stages of breast reconstruction at your 6 week follow-up appointment.**

### YOUR FIRST YEAR

- **Continue your breast massage, and practice monthly breast self exam.**
- **Continue healthy nutrition, fitness and sun protection.**
- **Your scars will continue to refine.** If they become raised, red or thickened, or appear to widen, contact our office. Early intervention is important to achieving well-healed scars. Scars are generally refined to fine incision lines one year after surgery.
- **A one-year post surgery follow-up is required.** However you may call our office at any time with your concerns or for needed follow-up.
- **If your breasts begin to develop an unusually hard feeling, or a highly rounded “squeezed” appearance, call us as soon as possible.** Early treatment is the best solution to capsular contracture. Breast massage is the most important form of early intervention.

**Your body will change with age.** The appearance of your breasts will change too. You may wish to have your implants replaced or to undergo revision surgery again in the future to help maintain your appearance throughout life. Contact our office with any of your questions or concerns, at any time.

**I have read and understand all of the above instructions. I understand that following these instructions is solely my responsibility. I understand that it is also my responsibility to ask my doctor and his or her staff any questions I have related to these instructions or about my procedure, health and healing.**

**These instructions are not meant to be comprehensive or all-inclusive, any additional instructions will be given to you by Dr. Butterfield as needed.**

**Financial Policy Regarding Revision and Complications**

Every plastic and reconstructive surgeon has a few patients who will require revision or have some complications requiring additional surgery. As you have been or will be told, one cannot guarantee a result. In cosmetic procedures there are certain problems that will happen statistically no matter how good the care or how careful the doctor and team. Examples of problems that may be encountered are bleeding or an unfavorable scar after a surgical procedure. In both of these cases, it may be necessary to return the patient to surgery, either on an emergency basis (as in the case with bleeding) or an elective basis (as in the case of scarring). It is our policy as a predetermined courtesy to our patients not to charge a surgeon's fee for complications or revisional surgery within 6 months from the original surgery date. We do, however, expect the patient to pay whatever other expenses arise as a result of treatment in hospital or outpatient settings. If the revisional surgery occurs in our office facility, the patient is responsible for the expense of the facility and anesthesia. Sometimes the patient will have insurance that will cover these revisions or complications. It depends upon the individual policy and how it is written. When a person does have insurance, the insurance company is billed for the surgeon's fee as well as the facility fees. All patient copays and deductibles will apply.

We hope that no complication arises and no revisional surgery is necessary in your case. However, no plastic surgeon can guarantee this to patients. It is important for the patient undergoing an elective surgical procedure to understand this financial policy. If you have any questions regarding this policy, the office staff would be happy to discuss it with you.

**Policy Regarding Employer Paperwork**

Many patients have forms that must be completed by Dr. Butterfield or our office to verify surgery dates, length of absence from work, work restrictions and expected return to work date. Many employers require multiple forms with ongoing updates. We understand our patients need to comply with their employer's human resource requirements. With regret the volume and complexity of the required paperwork makes it necessary for our office to charge a fee for this service. For those patients with human resource requirements each surgery will be assessed a \$20.00 fee for completion of all forms. All spouse paperwork will also be assessed a \$20.00 fee.

Please ensure all forms are in our office at least two weeks prior to your employers due date marked "Attention Nurse – Employer Forms". Include the address or fax number where all completed forms are to be returned, due date and your contact information should questions arise. The \$20.00 fee may be paid to the receptionist at the time of your pre operative appointment.

My signature below, indicates that I understand and agree to the above policies.

Signature \_\_\_\_\_ Date \_\_\_\_\_  
Witness \_\_\_\_\_